

Feedback Heating with Slow Jets in Cooling Flow Clusters

Noam Soker

Department of Physics, Technion–Israel Institute of Technology, Haifa 32000

`soker@physics.technion.ac.il`

and

Fabio Pizzolato

Department of Physics, Technion–Israel Institute of Technology, Haifa 32000

`fabio@physics.technion.ac.il`

ABSTRACT

We propose a scenario in which a large fraction, or even most, of the gas cooling to low temperatures of $T < 10^4$ K in cooling flow clusters, directly gains energy from the central black hole. Most of the cool gas is accelerated to non-relativistic high velocities, $v_j \simeq 10^3 - 10^4$ km s $^{-1}$, after flowing through, or close to, an accretion disk around the central black hole. A poorly collimated wind (or double not-well collimated opposite jets) is formed. According to the proposed scenario, this gas inflates some of the X-ray deficient bubbles, such that the average gas temperature inside these bubbles (cavities) in cooling flow clusters is $kT_b \lesssim 100$ keV. A large fraction of these bubbles will be very faint, or not detectable, in the radio. The bright rims of these weak smaller bubbles will appear as ripples. We suggest that the X-ray ripples observed in the Perseus cluster, for example, are not sound waves, but rather the rims of radio-faint weak bubbles which are only slightly hotter than their environment. This scenario is incorporated into the moderate cooling flow model; although not a necessary ingredient in that model, it brings it to better agreement with observations. In the moderate cooling flow model a cooling flow does exist, but the mass cooling rate is $\lesssim 10\%$ of that in old versions of cooling flow models.

Subject headings: galaxies: clusters: general — cooling flows — intergalactic medium — X-rays: galaxies: clusters

1. Introduction

Recent *Chandra* and *XMM-Newton* observations of cooling flow (CF) clusters of galaxies have failed to detect the large amounts of cool gas predicted by the old versions of the cooling flow (CF) model (e.g. Tamura et al. 2001; Peterson et al. 2003, 2004; Molendi & Pizzolato 2001), reducing by at least one order of magnitude the expected mass cooling rate to low temperatures. These results raise several questions. Do CFs occur at all? If they do, what is the actual mass cooling rate? What is the fate of the cool gas? In many clusters an unhindered radiative loss would lead to a sizeable cooling flow in few 10^8 yr: since this does not seem to happen, some reheating is required (e.g. Fabian 2003), what is the nature of this heating mechanism?

In this paper we adopt the framework of the moderate CF model to address the fate of the cool gas, and suggest that a large fraction, and in some cases most, of the cool gas is being ejected back to the ICM by the active galactic nucleus (AGN) operating at the central cD galaxy. The approach we adopt here differs from those adopted by most authors (e.g. Binney & Tabor 1995; Tucker & David 1997; Ciotti & Ostriker 2001; Binney 2004; Omma et al. 2004; Mathews et al. 2004; for more references see Peterson et al. 2004), in that we *do* consider a CF, moderate one though. We shall conclude that the best agreement with the available data occurs if the jets are not well collimated.

In § 2 we shall sketch the envisaged scenario, putting it in the appropriate framework of the moderate CF model. In § 3 we review some hints to slow jets. Among them, a key role is played by the hot bubbles observed with *Chandra* in several CF clusters. In § 4 we shall discuss the properties of slow and massive jets, and learn that the bubbles’ properties seem to require slow and dense jets, which imply mass cooling rate of $\gtrsim 10 M_\odot \text{ yr}^{-1}$ in large CF clusters. In other words, *the bubbles imply the presence of a CF*. We summarize in § 5.

2. The Proposed Scenario

We start this Section by recalling some important features of the moderate CF model (for more details, see Soker et al. 2001; Soker & David 2003), as well as some other relevant ingredients for our proposed model.

The moderate CF model, which was proposed before the new results from *Chandra* and *XMM-Newton* (hence we avoid referring to old version of the CF model as *standard*), is different from many earlier proposed processes whose aim is to prevent CF in clusters of galaxies altogether (Soker et al. 2001; Soker & David 2003). The main ingredient of the moderate CF model is that the effective age, i.e. the time elapsed since the last major

disturbance of the intra-cluster medium (ICM) inside the cooling radius $r_c \simeq 100$ kpc, is much shorter than the cluster age (e.g. Binney & Tabor 1995; Binney 2004; Soker et al. 2001). The cooling radius is defined as the place where the radiative cooling time equals the cluster age. Originally, the heating in the moderate CF model was proposed to be intermittent (Soker et al. 2001), but the basic idea can hold for a steady heating, or heating in short intervals (Binney 2004).

In the moderate CF model most of the gas within the CF resides in the hottest phase, which is prevented from cooling continuously and attaining a steady-state configuration by being reheated (Soker & David 2003; Kaiser & Binney 2004). This results in a mass cooling rate that decreases with decreasing temperature, with a much lower mass cooling rate at the lowest temperatures. The limit on the cooling rate below a temperature $T_{\min}$ inferred from X-ray observations is $< 20\%$ of the mass cooling rates cited in the past (e.g. Fabian et al. 2002a; Molendi & Pizzolato 2001; Peterson et al. 2003). In some cases, though, cooling to low temperatures is observed, as in the CF clusters Abell 2597 (Morris & Fabian 2004) and Abell 2029 (Clarke et al. 2004). In Abell 2597 both extreme-UV and X-ray observations indicate a mass cooling rate of $\sim 50 M_{\odot} \text{ yr}^{-1}$, which is ~ 0.2 of the value quoted in the past based on *ROSAT* X-ray observations (see the discussion in Morris & Fabian 2004). Some fraction of the gas may cool to lower temperatures by heat transfer to optical-emitting gas, reducing further the X-ray emission from gas residing at temperatures of $T \lesssim T_{\min} \simeq 1$ keV; the energy transfer can be via mixing (e.g. Oegerle et al. 2001; Fabian et al. 2001, 2002a; Johnstone et al. 2002; Bayer-Kim et al. 2002), and/or heat conduction (Soker et al. 2004; Soker 2004b). Intermittent heating, which is part of the moderate CF model, probably occurs by AGN activity. In a recent paper McNamara et al. (2004a) found a huge deficient X-ray bubble pair in the cluster MS 0735.6+7421, which suggests that an intermittent heating over a large time scale might play a role. In the moderate cluster CF model, the agreement between star formation rate and mass cooling rate can be quite good. Wise et al. (2004) and McNamara et al. (2004b), for example, find the cooling rate within $r \simeq 30$ kpc of the CF cluster A 1068 to be about equal to the star formation rate there ($20 - 70 M_{\odot} \text{ yr}^{-1}$).

Another plausible ingredient of the moderate CF model is that the entire inner CF region supplies the cold gas accreted to the black hole (Pizzolato & Soker 2004). Namely, the feedback between heating and cooling occurs with the entire cool inner region, $r \lesssim 5-30$ kpc, in what we term a *cold-feedback model*. In the proposed scenario (Pizzolato & Soker 2004) non-linear over-dense blobs of gas, $\delta\rho/\rho_a \gtrsim 2$, i.e., $\rho/\rho_a \gtrsim 3$, in this inner region cool on short time scales such that they are removed from the ICM before the next major AGN heating event in their region. Some of these blobs cool and sink toward the central black hole, while others may form stars and cold molecular clouds. This mechanism can work on the condition that the blobs have a small angular momentum, or they would be prevented

from accreting onto the central black hole and fuelling its activity. This issue has been discussed in a separate paper (Pizzolato & Soker 2004): here we just summarize the main conclusions. The cold blobs may stem directly from ICM disturbances driven by an earlier AGN activity, but also from galaxies mass-stripping (Soker et al. 1991). Since the galaxies do not have an ordered bulk motion, the mass stripped from them is also unlikely to conspire and organize in an ordered flow with high net angular momentum. Therefore, if a circular flow like a disc forms, it cannot be very large: say, $\sim 10^2$ pc as in M87 (Harms et al. 1994; Ford et al. 1994). The orbit of a dense blob subjected to gravity and the friction drag force has a circularization radius whose size is in broad agreement with this value (Pizzolato & Soker 2004). Besides, in the cold feedback model the cold gas is expected to accrete from regions not too far from the center (few tens kpc at most), so the accreting flow should not have a large angular momentum from the outset. Therefore, the blobs’ angular momentum is not high enough to prevent them from accreting in a time scale shorter than or comparable to the cooling time, which keeps running the cooling/accretion feedback loop.

In the present paper we suggest that a large fraction, and in some cases most, of the accreting cool gas is being ejected back to the ICM by the AGN. This idea solves some problems related to the questions posed above. In particular, the slower and more massive collimated wind (namely, a poorly collimated jet pair; hereafter referred to simply as jets), deposits its energy in the inner region (Binney 2004), as required by recent observations that limit the degree of mixing (Böhringer et al. 2004), and can carry more energy than inferred from radio observations (Binney 2004). In the present paper we extend this idea of slow and massive not-well collimated jets, and incorporate it to be an important ingredient in the duty-cycle of the moderate CF model. The proposed scenario is fundamentally different from the scenario of Nulsen (2004) or that of Binney (2004) (see also Omma & Binney 2004). In those papers, most, or all, of the mass which is accreted to the central black hole comes from the hot phase, $T \simeq 10^7$ K. Hence, the mass that is ejected in the jets is small compared to the mass assumed to be cooling to $T \lesssim 10^4$ K in the moderate CF model. In the proposed scenario, on the other hand, we address the question of the fate of the cold gas, arguing that non negligible fraction of it is ejected back to the ICM at high non-relativistic velocities. In those papers the duty-cycle, or feedback process, is determined by energy considerations alone, while in the present scenario the mass is also a factor in the duty-cycle.

Our proposed scenario is also different from the circulation flow proposed by Mathews et al. (2004); in their scenario the gas does not cool below X-ray emission temperature. This is a significant difference, as basically they don’t consider the presence of a CF to low temperatures ($T < 10^4$ K), while we do. In common with their model, we ascribe significance to mass as well as energy transport in both inward and outward direction.

The cool ICM flows to the center, and if it has an angular momentum, it may form an accretion disc about the central black hole. A standard geometrically thin, optically thick accretion discs (à la Shakura-Sunyaev) truncated at the last marginally stable orbit around the black hole has a radiative efficiency $\eta \simeq 10\%$, corresponding to a bolometric luminosity

$$L_{\text{accr}} \simeq 5.7 \times 10^{45} \text{erg s}^{-1} \left(\frac{\eta}{0.1} \right) \left(\frac{\dot{M}}{M_{\odot} \text{ yr}^{-1}} \right), \quad (1)$$

Such bright discs are not common, which forces us to assume that the disc’s radiative efficiency is low. The disc may be truncated at several Schwarzschild radii from the hole (e.g. it may be evaporated by magnetic fields), which reduces η in Equation (1) by orders of magnitude. Alternatively, the disc may be radiatively inefficient, i.e. it does not radiate most of the energy converted by the viscous dissipation: such is an *ADAF* or an *ADIOS* (Narayan & Yi 1994; Blandford & Begelman 1999). Omma et al. (2004) assume an *ADIOS* flow in the framework of their simulations. In these kinds of accretion flow, most of the incoming flux is gravitationally unbound to the black hole, and can therefore be pushed back as an outflow, or a wind. A possible issue is that *ADAFs* cannot exist for high accretion rates (Quataert et al. 1999). There is the possibility, however, that the disc is composite, with the *ADAF* component confined in the inner part, the outer being an ordinary accretion disc (e.g. Quataert et al. 1999). Discs like these may exist even for high accretion rates, and may also support the required outflow.

In the next section we start by reviewing some hints that the outflowing jets are slow. It seems as if the properties of bubbles require slow and dense jets, which imply mass cooling rate of $\gtrsim 10 M_{\odot} \text{ yr}^{-1}$ in large CF clusters. Namely, *the bubbles imply the presence of a CF*. We then estimate the relevant parameters, and summarize by listing some predictions of the proposed model.

3. The Temperature of Hot Bubbles

We list below some pieces of evidence that some bubbles are inflated mainly by non-relativistic jets, i.e., the temperature of the gas inside some bubbles is $T_b \lesssim 100 \text{ keV}$. A population of radio emitting relativistic electrons may exist due to the contribution from a relativistic jet as well. In other cases, like in the large bubbles in M87, the inflating jets are relativistic (Forman et al. 2003), and the non-relativistic component is lacking.

1. In most clusters only a lower limit determination of the hot bubbles temperature is possible (Blanton 2004). Some examples are $T_b > 15 \text{ keV}$ for Hydra A (Nulsen et al.

2002), $T_b > 11$ keV for Perseus (Schmidt et al. 2002), and $T_b > 20$ keV for Abell 2052 (Blanton et al. 2003). We are aware only of the case of MKW 3S where a bubble temperature was measured (Mazzotta et al. 2002). In this cluster a surface brightness depression located at a distance of $r_b \simeq 100$ kpc from the cluster center has a temperature $T_b \simeq 6 - 10$ keV (Mazzotta et al. 2002). Both in Hydra A (David et al. 2001) and Abell 2052 (Blanton et al. 2001) the pressure drops by a factor of $f_p \simeq 0.1$ from the inner region to $r \simeq 100$ kpc. For an adiabatic expansion with a constant ratio of the bubble to external pressure, the temperature inside the bubble drops by a factor of $f_t = f_p^{2/5} \simeq 0.4$. If we assume that the bubble in MKW 3S was inflated in the inner region, and then was buoyantly rising and adiabatically expanding, then its initial temperature was $T_{b0} \simeq 15 - 25$ keV. The preliminary results of McNamara et al. (2004a) show that the large two X-ray deficient bubbles in the cluster MS 0735.6+7421 are only slightly hotter than their environment, strengthening the results above. A word of caution about these measurements is in order, however. It is extremely difficult to measure the temperature of the material in the ICM bubble with the current data. The geometry of the cavities is never known well enough to allow an accurate deprojection; besides it is unknown whether the emission stems from the body of the cavity or from its rims. If we were to adopt a more conservative viewpoint, we could say that the current temperature measurements are *consistent* with the view that these blobs are filled with hot plasma.

2. The pressure in the bubbles inferred from the radio emission and the assumption of equipartition between magnetic and relativistic particles is lower than the ambient pressure (Blanton 2004). This hints (but does not *prove*), that the bubbles are not entirely relativistic, because the bubbles are in pressure equilibrium with their environment.
3. Some AGNs have been observed to blow non-relativistic winds, with speeds down to $24,000 \text{ km s}^{-1}$ (PDS 456: Reeves et al. 2003, PG1211+143: Pounds et al. 2003; however, see McKernan et al. 2004 for some possible problems). The identification of an even slower wind ($\sim 1,000 \text{ km s}^{-1}$) in the *LINER* NGC 1097 is more problematic (Storchi-Bergmann et al. 2003, and references therein).

On the theoretical side, Binney (2004) lists some arguments in support of a slow outflow. According to Binney (2004), in many cases the relativistic jet carries a small fraction of the mass and energy in the outflow, and in some systems no relativistic jet is present even when slower outflow occurs. In their bubble-inflation simulations, Omma & Binney (2004) take the wind speed to be in the range $2.8 - 4 \times 10^4 \text{ km s}^{-1}$, with a mass loss rate of $1 M_\odot \text{ yr}^{-1}$, and Omma et al. (2004) take $v_j = 10,000 \text{ km s}^{-1}$ and a mass loss rate into the jet of $2 M_\odot \text{ yr}^{-1}$. In many models for quasars the velocity of the wind blown by the disk is in the range $\sim 10^3 - 3 \times 10^4 \text{ km s}^{-1}$ (Elvis 2000;

Crenshaw et al. 2003, and references therein). There are other models which predict slow ($\lesssim 10^4 \text{ km s}^{-1}$) winds, with the most recent one being the one by Begelman & Celotti (2004). In the model proposed by Nicastro (2000), the wind velocity decreases from $\sim 20,000 \text{ km s}^{-1}$ to $\sim 1000 \text{ km s}^{-1}$ as the mass accretion rate increases; in this model the high accretion rate expected in CF clusters may result in slow winds. In Seyfert galaxies, mass loss rates from the black hole vicinity as large as $\simeq 1 M_{\odot} \text{ yr}^{-1}$ have been inferred (Crenshaw et al. 2003). The outflow of the absorbing gas starts at distances as small as $\sim 0.01 \text{ pc}$ from the black hole, indicating origin from the accretion disk. The possibility of slow jets from the nucleus (3C 317) of the cooling flow cluster A2052 was raised recently by Venturi et al. (2004).

The slow wind velocities suggested above leads to bubble temperatures of $kT_B \lesssim 100 \text{ keV}$. Using the expression given by Castor et al. (1975) for the expansion of an interstellar bubble (Bicknell & Begelman 1996, also applied this expression to study the bubble formed by the radio jet in M87), the temperature inside the bubble is

$$kT_B \simeq 0.15 \mu m_H v_j^2 = 100 \left(\frac{v_j}{10^4 \text{ km s}^{-1}} \right)^2 \text{ keV}, \quad (2)$$

where v_j is the jet velocity, and μm_H the mean mass per particle. This temperature implies a low density in the bubble, assuming pressure equilibrium with the ICM. Even for $v_j = 4000 \text{ km s}^{-1}$, the bubble temperature is $\gtrsim 5$ times higher than the ambient temperature in the inner regions of CF clusters (typically $\simeq 1 - 3 \text{ keV}$). This means a bubble density of $n \lesssim 0.2$ times the ambient density, and emissivity $n^2 \lesssim 0.04$ times that of the ambient medium. This low emissivity cannot be detected by present X-ray telescopes.

If, as assumed here following Binney (2004), the AGN blows gas with a spectrum of velocities, then the bubble is multi-phase. In some cases the fastest blown gas has relativistic speeds, and the bubble is detectable in the radio. The formation process of the bubble is likely to lead to some internal motion within the bubble. This internal motion determines the mixing of the different gas-phases at later times, and the magnetic field topology and evolution determines local heat conduction. The evolution at late times is hard to predict, and probably very hard to simulate as well. We do expect that the phases will be mixed among themselves, and later on with the ICM, hence heating the ICM. As with other bubbles models, the bubbles themselves heat the ICM by doing work on the ICM, and by lifting cool ICM medium from the center.

We stress that in the present model slow winds do *not* exclude the occurrence of fast, relativistic jets. Indeed, many of the best examples of ICM cavities are clearly associated with radio lobes (e.g. Hydra-A: McNamara et al. 2000; Perseus: Fabian et al. 2002b). Slow winds and relativistic jets may coexist: the former carry most of the energy (Binney 2004),

and the latter fuel of the synchrotron emission from the relativistic electrons in the X-ray cavities.

4. Jet Properties

4.1. The Mass Flow Rate into the Jets

Combining the estimated energy of bubbles (or bubble pairs) with the typical non-relativistic velocity assumed in the previous Section yields the mass ejection rate. Birzan et al. (2004) give the observed mechanical luminosity of 16 clusters. The jets' kinetic luminosity will be equal to several times the power L_{mech} calculated by them from the bubbles values of PV , where P is the pressure and V the bubbles' volume. Most luminous clusters reside, therefore, in the range $\dot{E}_j \simeq 0.3 - 3 \times 10^{44} \text{erg s}^{-1}$, with possible higher values if the pressure inside bubbles is higher than in their surrounding (Birzan et al. 2004), or if some energy is in shocks formed by the inflated bubbles (Forman et al. 2003; McNamara et al. 2004a). Numerical simulation require the power to be on the upper side $\dot{E}_j \simeq 5 \times 10^{44} \text{erg s}^{-1}$ (e.g. Dalla Vecchia et al. 2004; Omma & Binney 2004). Also, the power of jets in AGN can reach a power of $\dot{E}_{j, \text{max}} \simeq 10^{47} \text{erg s}^{-1}$ (e.g. Rawlings & Saunders 1991). We therefore scale the averaged energy injected by the AGN in CF clusters with values higher than the energy directly observed (Binney 2004), and take it to be $5 \times 10^{44} \text{erg s}^{-1}$. Scaling with typical values and using Equation (2), the mass loss rate into the jets is

$$\dot{M}_j \simeq 15 \left(\frac{\dot{E}_j}{5 \times 10^{44} \text{erg s}^{-1}} \right) \left(\frac{kT_b}{100 \text{ keV}} \right)^{-1} M_{\odot} \text{ yr}^{-1}. \quad (3)$$

In MKW 3S the temperature inside the bubble T_b has been estimated (Mazzotta et al. 2002), and we take the value calculated in the previous section for the initial value of the temperature, $T_{b0} \simeq 20 \text{ keV}$, and take also $\dot{E} \gtrsim 10^{44} \text{erg s}^{-1}$ found by Birzan et al. (2004). From these we find $\dot{M}_j \gtrsim 15 M_{\odot} \text{ yr}^{-1}$. This is more than the X-ray-inferred mass cooling rate of $\dot{M}_{\text{cool}} < 2 M_{\odot} \text{ yr}^{-1}$ Birzan et al. (2004). However, in MKW 3S the bubbles are not so prominent as in other clusters, and may represent a relatively old ejection event.

4.2. Jet Propagation

Based on § 3 of Soker (2004a), we note the following properties in regards to the heavy jets. It is assumed that the slow collimated wind (or not-well collimated jet) has a wide opening angle, measured from the symmetry axis of the jet, of $\alpha \simeq 1$. The expansion

velocity of the jet's head v_h is given by the following expression as long as $v_h \ll v_j$:

$$v_h \simeq \left[\frac{\dot{E}_j}{\pi(1 - \cos \alpha) z^2 v_j \rho_c} \right]^{1/2} = 1200 \left(\frac{\dot{E}_j}{5 \times 10^{44} \text{erg s}^{-1}} \right)^{1/2} \left(\frac{v_j}{10^4 \text{ km s}^{-1}} \right)^{-1/2} \\ \times \left(\frac{1 - \cos \alpha}{0.5} \right)^{-1/2} \left(\frac{\rho_c}{10^{-25} \text{g cm}^{-3}} \right)^{-1/2} \left(\frac{z}{5 \text{ kpc}} \right)^{-1} \text{ km s}^{-1}, \quad (4)$$

where ρ_c is the ambient density, and the z the distance of the jet's head from its source, measured along the jet's symmetry axis. Note that the jet is scaled with an opening angle of $\alpha = 60^\circ$ (from its symmetry axis; the full opening angle is 120°). For these parameters the jet becomes subsonic in the ICM at a distance of ~ 10 kpc. The subsonic jet expands to the side, ensuring the formation of a large bubble. A large opening angle, although not a necessary condition to inflate a bubble, facilitates the formation of a large bubble close to the center.

A bubble can be formed before the jet's head becomes subsonic if the shocked material in the jet expands faster than the jet's head (Soker 2004a). The condition on the opening angle of the jet for that to occur is given by Equation (14) of Soker (2004a). We change the variable in that equation as follows. From Equation (4), the distance of the jet's head as function of time is given by

$$z \simeq 3.4 \left(\frac{t}{10^6 \text{ yr}} \right)^{1/2} \text{ kpc}. \quad (5)$$

We substitute the value of the time from the last equation into Equation (14) of Soker (2004a) to derive the condition to inflate a bubble via the mechanism discussed in Soker (2004a)

$$\alpha \gtrsim 50^\circ \left(\frac{\dot{E}_j}{5 \times 10^{44} \text{erg s}^{-1}} \right)^{3/10} \left(\frac{v_j}{10^4 \text{ km s}^{-1}} \right)^{-1/2} \left(\frac{\rho_c}{10^{-25} \text{g cm}^{-3}} \right)^{-3/10} \left(\frac{z}{10 \text{ kpc}} \right)^{-1/5}. \quad (6)$$

From Equations (4) and (6) we see that the large opening angle of the jet (hence termed here a not-well collimated jet) facilitates bubble formation (Soker 2004a). In the proposed scenario a bubble is formed at a distance of $z \simeq 10$ kpc from the source of the jet. The large opening angle assumed here is different from the heavy slow jet simulation of Omma et al. (2004).

The large opening angle implies that the jet, when expands outward, interacts with a substantial fraction of the ICM in its vicinity. The jet, and the bubble it forms, pushes outward ICM medium which is cooler and denser than the ICM the bubble reaches at later time. Such a wide open angle flow might account for dense shells observed around some bubbles, e.g., in Abell 2052 (Blanton et al. 2001).

With our proposed scenario, where many small and weak bubbles which are only slightly hotter than their environment exist, we turn to the deep X-ray image of the Perseus CF cluster (Fabian et al. 2003). Many bright arcs, which are termed ripples, are seen in this cluster. They were interpreted by Fabian et al. (2003) as sound waves. However, we suggest that these are actually the bright rims of a stack of weak flattened bubbles accumulated in the past, which have been flattened by the resistance of the environment ICM they push through. This is supported by noting the following: (1) The rims of the two inner X-ray deficient strong bubbles, and the rim of the outer strong bubble, look like the X-ray ripples, both in shape and size; only that these rims are brighter than the ripples. (2) Some ripples, when considered as geometric spherical arcs, have their center off from the cluster center. (3) While sound waves are expected to expand to all directions, some ripples are very short. One possible prediction of our proposal that the ripples are actually rims of weak bubbles, is that very weak radio emission will be detected between some ripples, similar to, but much weaker than, the radio emission in X-ray deficient bubbles.

On the other extreme, a slow jet can be well collimated, hence propagates along a narrow cone into the ICM. Let β be defined such that the jet expands into a solid angle of $\Omega_j = 4\pi\beta$. For $\alpha = 12^\circ$, for example, $\beta = 0.01$. By neglecting the magnetic pressure inside the jet and relativistic effects, hence $\dot{E}_j = \dot{M}_j v_j^2 / 2$, the speed of the jet's head is determined by pressure equilibrium on its two sides. An approximate relation is obtained if we consider only ram pressures, $\rho_j (v_j - v_h)^2 = \rho_a v_h^2$, where ρ_a is the ICM density, $\rho_j = \dot{M}_j / (4\pi\beta r^2 v_j)$, is the density inside the jet, v_j is the speed of the gas inside the jet, v_h the speed of the jet's head, $\dot{M}_j$ is the mass loss rate into one jet, and r is the distance of the jet's head from its origin (e.g. Krause 2003). For $v_j \gg v_h$, the jet's head propagation speed is given by

$$v_h = \frac{v_j}{(\rho_a/\rho_j)^{1/2} + 1} \simeq 600 \left(\frac{\dot{E}_j}{5 \times 10^{43} \text{ erg s}^{-1}} \right)^{1/2} \left(\frac{v_j}{10^5 \text{ km s}^{-1}} \right)^{-1/2} \\ \times \left(\frac{\beta}{0.01} \right)^{-1/2} \left(\frac{n_a}{0.1 \text{ cm}^{-3}} \right)^{-1/2} \left(\frac{r}{5 \text{ kpc}} \right)^{-1} \text{ km s}^{-1}. \quad (7)$$

We scaled the power of one jet according to that of Hydra A as given by Bîrzan et al. (2004), the total ambient number density is scaled with that of Hydra A as given by David et al. (2001), and we scaled with a very fast jet of $v_j = 0.33c$.

The last equation shows that jets like those in Hydra A, if they are very fast, become subsonic at $\sim 5 \text{ kpc}$ from their origin. This explains the observations that the jets in Hydra A turn into lobes. But what is the reason for the sharp transition from a well collimated jet to radio lobes in Hydra A? We note from the last equation that a slow jet, even if weak, might stay supersonic to large distances. For example, for a jet with $\dot{E} = 10^{43} \text{ erg s}^{-1}$ and

$v_j = 3000 \text{ km s}^{-1}$ we find $(\rho_a/\rho_j)^{1/2} \simeq 2$, and the exact solution of the last equation is $v_h \simeq 1000 \text{ km s}^{-1}$, at a distance of $r = 5 \text{ kpc}$. We propose, therefore, that the current jets in Hydra A were preceded by collimated slow and dense jets which opened a tunnel to a distance of $r \sim 5 \text{ kpc}$, through which the current jets are expanding almost undisturbed. When the current jets leave these tunnel, they interact with the ICM, become subsonic, and lose their collimation. A slow jet with these parameters ($\dot{E} = 10^{43} \text{ erg s}^{-1}$ and $v_j = 3000 \text{ km s}^{-1}$) has a mass loss rate of $3.5 M_\odot \text{ yr}^{-1}$, namely, the two proposed slow jets blow $\sim 7 M_\odot \text{ yr}^{-1}$ back to the ICM. To propagate to a distance of $\sim 5 \text{ kpc}$, the slow jets were active for $\text{few} \times 10^6 \text{ yr}$. Hence, in total, they blew a non negligible mass back to the ICM.

4.3. Spectral signature of the outflow

What are the observational characteristics of the slow poorly collimated outflow? Usually, the winds blown by AGNs are revealed by their absorption of the underlying continuum in the UV or X-rays (e.g. PG 1211+143: Pounds et al. 2003; PDS 456: Reeves et al. 2003; NGC 1097: Storchi-Bergmann et al. 2003; see also Crenshaw et al. 2003 for a recent review). Since the outflows we have dealt with in this paper are not very hot at their outset ($10^5 - 10^6 \text{ K}$), also their ionization degree should allow detection in the UV or in X-rays, as in ordinary AGNs. Since the wind remains relatively cold on rather small, sub-kpc scales, the actual detectability of the wind may be complicated by other factors, like the absorption by intervening clouds. The slow outflow is active during a small fraction (10 – 30%) of the time. Hence, most clusters will not show any signature of the slow wind.

5. Summary

The purpose of the current paper is to propose a plausible new ingredient for the moderate cooling flow (CF) model. We propose (speculate) that a large fraction, or even most, of the mass that cools to low temperatures is ejected back to the hot ICM via an accretion disk around the central black hole. In the moderate CF model the effective age of the CF is much shorter than the cluster age, because of intermittent heating; most likely via AGN activity; only in the very inner regions the flow is in a steady state phase; cooling to low temperatures ($T < 10^4 \text{ K}$) occurs at a rate much lower than in older versions of CF models (see § 1).

This proposed scenario is based on several observations and theoretical considerations (§ 3) that hint either at bubbles temperatures of $T_b \lesssim 100 \text{ keV}$, or at the possibility that

AGN can blow relatively slow winds (jets), $1000 \lesssim v_j \lesssim 10,000 \text{ km s}^{-1}$. With the lower energy per ejected unit mass (Equation (2)), the outflowing mass required to account for the energy in the bubbles may comprise a significant fraction of the mass cooling to low temperature (Equation (3)). In the proposed scenario, the average power of the AGN is much higher than that inferred from radio emission, or even from energy content of the large bubbles. Some mass will be blown at $v_j \simeq 3000 - 5000 \text{ km s}^{-1}$, forming small bubbles, with density not much smaller than their surroundings. Bîrzan et al. (2004) find that the energy associated with X-ray deficient bubbles does not generally explain the low mass cooling rate, unless the bubbles probe only a small fraction of the total kinetic energy. The proposed speculative scenario accounts for this extra energy.

In the proposed scenario most of the cooling mass stays in the inner region $r \lesssim 10 - 50 \text{ kpc}$ of the CF cluster, as most bubbles do not buoy to large radii. This is compatible with the implication that there is no vigorous mixing in CF clusters (Böhringer et al. 2004). We stress that there is no wind of the ICM, simply we proposed that some fraction of the jets (or collimated outflow), which inflate some of the bubbles, is blown at a relatively low speed, and contains significant fraction of the mass cooling to temperatures of $\lesssim 10^4 \text{ K}$. This ingredient of the moderate CF model makes better the agreement of the model with observations. However, this ingredient is not necessary for the moderate CF model.

The proposed scenario predicts the following.

1. As being an ingredient in the moderate CF model, it requires that some mass cools to low temperatures, but at a moderate average rates $\dot{M}_{\text{cool}} \simeq 1 - 50 \text{ } M_{\odot} \text{ yr}^{-1}$.
2. The temperatures of the gas in large bubbles is, in most cases, $T_b \lesssim 100 \text{ keV}$.
3. In many clusters small bubbles exist, which are only slightly hotter than their surroundings. Such bubbles may reveal themselves as patchy low-X-ray emitting regions, such as those observed in M87-Virgo (see the X-ray image by Young et al. 2002).
4. Weak radio emission between X-ray ripples, as the ripples discovered by Fabian et al. (2003) in the Perseus cluster. We recognize, however, that these observations may be difficult, chiefly on account of confusing projection effects.
5. Sub-relativistic, $v_j \lesssim 10,000 \text{ km s}^{-1}$, flow from relatively extended region around the central black hole of CF clusters exists from time to time (or presently, for 10 – 30% of the cooling-flow clusters).
6. At least in some favourable cases, we do expect to detect such outflows thanks to their absorption in the UV or X-ray spectrum, as observed in the winds blown by other AGNs.

The first three predictions can be tested with present X-ray telescopes, We predict that deep X-ray observations of the inner regions of CF clusters will reveal: (1) emission from gas cooling below ~ 1 keV, as in Abell 2597 (Morris & Fabian 2004) and Abell 2029 (Clarke et al. 2004); (2) some X-ray emission from some bubbles, implying their density is not much lower than their ambient density; and (3) small bubbles scattered around, only slightly hotter than their surroundings.

We thank Orkan M. Umurhan for comments on the original manuscript, and Shai Kaspi for useful discussion. We also thank an anonymous referee for insightful comments. This research was partially supported by a grant from the Israel Science Foundation. F. Pizzolato was supported also by grant No. 2002111 from the United States-Israel Binational Foundation, Jerusalem, Israel. He is currently supported by a Fine Fellowship.

REFERENCES

- Bîrzan, L., Rafferty, D. A., McNamara, B. R., Wise, M. W., & Nulsen, P. E. J. 2004, *ApJ*, 607, 800
- Böhringer, H., Matsushita, K., Churazov, E., Finoguenov, A., & Ikebe, Y. 2004, *A&A*, 416, L21
- Bayer-Kim, C. M., Crawford, C. S., Allen, S. W., Edge, A. C., & Fabian, A. C. 2002, *MNRAS*, 337, 938
- Begelman, M. C. & Celotti, A. 2004, *MNRAS*, 352, L45
- Bicknell, G. V. & Begelman, M. C. 1996, *ApJ*, 467, 597
- Binney, J. 2004, in *The Riddle of Cooling Flows in Galaxies and Clusters of Galaxies*, published electronically at <http://www.astro.virginia.edu/coolflow/proc.php> (astro-ph/0310222)
- Binney, J. & Tabor, G. 1995, *MNRAS*, 276, 663
- Blandford, R. D. & Begelman, M. C. 1999, *MNRAS*, 303, L1
- Blanton, E. L. 2004, in *The Riddle of Cooling Flows in Galaxies and Clusters of Galaxies*, published electronically at <http://www.astro.virginia.edu/coolflow/proc.php> (astro-ph/0402342)

- Blanton, E. L., Sarazin, C. L., & McNamara, B. R. 2003, *ApJ*, 585, 227
- Blanton, E. L., Sarazin, C. L., McNamara, B. R., & Wise, M. W. 2001, *ApJ*, 558, L15
- Castor, J., McCray, R., & Weaver, R. 1975, *ApJ*, 200, L107
- Ciotti, L. & Ostriker, J. P. 2001, *ApJ*, 551, 131
- Clarke, T. E., Blanton, E. L., & Sarazin, C. L. 2004, *astro-ph/0408068*, *ApJ*, in press
- Crenshaw, D. M., Kraemer, S. B., & George, I. M. 2003, *ARA&A*, 41, 117
- Dalla Vecchia, C., Bower, R. G., Theuns, T., Balogh, M. L., Mazzotta, P., & Frenk, C. S. 2004, *MNRAS*, 507
- David, L. P., Nulsen, P. E. J., McNamara, B. R., Forman, W., Jones, C., Ponman, T., Robertson, B., & Wise, M. 2001, *ApJ*, 557, 546
- Elvis, M. 2000, *ApJ*, 545, 63
- Fabian, A. C. 2003, *MNRAS*, 344, L27
- Fabian, A. C., Allen, S. W., Crawford, C. S., Johnstone, R. M., Morris, R. G., Sanders, J. S., & Schmidt, R. W. 2002a, *MNRAS*, 332, L50
- Fabian, A. C., Celotti, A., Blundell, K. M., Kassim, N. E., & Perley, R. A. 2002b, *MNRAS*, 331, 369
- Fabian, A. C., Mushotzky, R. F., Nulsen, P. E. J., & Peterson, J. R. 2001, *MNRAS*, 321, L20
- Fabian, A. C., Sanders, J. S., Crawford, C. S., Conselice, C. J., Gallagher, J. S., & Wyse, R. F. G. 2003, *MNRAS*, 344, L48
- Ford, H. C., Harms, R. J., Tsvetanov, Z. I., Hartig, G. F., Dressel, L. L., Kriss, G. A., Bohlin, R. C., Davidsen, A. F., Margon, B., & Kochhar, A. K. 1994, *ApJ*, 435, L27
- Forman, W., Nulsen, P., Heinz, S., Owen, F., Eilek, J., Vikhlinin, A., Markevitch, M., Kraft, R., Churazov, E., & Jones, C. 2003, (*astro-ph/0312576*)
- Harms, R. J., Ford, H. C., Tsvetanov, Z. I., Hartig, G. F., Dressel, L. L., Kriss, G. A., Bohlin, R., Davidsen, A. F., Margon, B., & Kochhar, A. K. 1994, *ApJ*, 435, L35
- Johnstone, R. M., Allen, S. W., Fabian, A. C., & Sanders, J. S. 2002, *MNRAS*, 336, 299

- Kaiser, C. R. & Binney, J. J. 2004, in *The Riddle of Cooling Flows in Galaxies and Clusters of Galaxies*, published electronically at <http://www.astro.virginia.edu/coolflow/proc.php>
- Krause, M. 2003, *A&A*, 398, 113
- Mathews, W. G., Brighenti, F., & Buote, D. A. 2004, *astro-ph/0407367*, (*ApJ*, in press)
- Mazzotta, P., Kaastra, J. S., Paerels, F. B., Ferrigno, C., Colafrancesco, S., Mewe, R., & Forman, W. R. 2002, *ApJ*, 567, L37
- McKernan, B., Yaqoob, T., & Reynolds, C. S. 2004, (*astro-ph/0408506*)
- McNamara, B. R., Nulsen, P. E. J., Wise, M. W., Bîrzan, L., Rafferty, D. A., Carilli, C. L., Blanton, E. L., & Sarazin, C. L. 2004a, Presented at the 35th COSPAR Scientific Assembly, Paris, July 19-20, 2004.
- McNamara, B. R., Wise, M., Nulsen, P. E. J., David, L. P., Sarazin, C. L., Bautz, M., Markevitch, M., Vikhlinin, A., Forman, W. R., Jones, C., & Harris, D. E. 2000, *ApJ*, 534, L135
- McNamara, B. R., Wise, M. W., & Murray, S. S. 2004b, *ApJ*, 601, 173
- Molendi, S. & Pizzolato, F. 2001, *ApJ*, 560, 194
- Morris, G. & Fabian, A. C. 2004, Presented at the 35th COSPAR Scientific Assembly, Paris, July 19-20, 2004.
- Narayan, R. & Yi, I. 1994, *ApJ*, 428, L13
- Nicastro, F. 2000, *ApJ*, 530, L65
- Nulsen, P. 2004, in *The Riddle of Cooling Flows in Galaxies and Clusters of Galaxies*, published electronically at <http://www.astro.virginia.edu/coolflow/proc.php> (*astro-ph/0310195*)
- Nulsen, P. E. J., David, L. P., McNamara, B. R., Jones, C., Forman, W. R., & Wise, M. 2002, *ApJ*, 568, 163
- Oegerle, W. R., Cowie, L., Davidsen, A., Hu, E., Hutchings, J., Murphy, E., Sembach, K., & Woodgate, B. 2001, *ApJ*, 560, 187
- Omma, H. & Binney, J. 2004, *MNRAS*, 350, L13

- Omma, H., Binney, J., Bryan, G., & Slyz, A. 2004, MNRAS, 348, 1105
- Peterson, J. R., Kahn, S. M., Paerels, F. B. S., Kaastra, J. S., Tamura, T., Bleeker, J. A. M., Ferrigno, C., & Jernigan, J. G. 2003, ApJ, 590, 207
- Peterson, J. R., Kahn, S. M., Paerels, F. B. S., Kaastra, J. S., Tamura, T., Bleeker, J. A. M., Ferrigno, C., & Jernigan, J. G. 2004, in *The Riddle of Cooling Flows in Galaxies and Clusters of Galaxies*, published electronically at <http://www.astro.virginia.edu/coolflow/proc.php>
- Pizzolato, F. & Soker, N. 2004, submitted to the MNRAS
- Pounds, K. A., Reeves, J. N., King, A. R., Page, K. L., O’Brien, P. T., & Turner, M. J. L. 2003, MNRAS, 345, 705
- Quataert, E., di Matteo, T., Narayan, R., & Ho, L. C. 1999, ApJ, 525, L89
- Rawlings, S. & Saunders, R. 1991, Nature, 349, 138
- Reeves, J. N., O’Brien, P. T., & Ward, M. J. 2003, ApJ, 593, L65
- Schmidt, R. W., Fabian, A. C., & Sanders, J. S. 2002, MNRAS, 337, 71
- Soker, N. 2004a, A&A, 414, 943
- . 2004b, MNRAS, 350, 1015
- Soker, N., Blanton, E. L., & Sarazin, C. L. 2004, A&A, 422, 445
- Soker, N., Bregman, J. N., & Sarazin, C. L. 1991, 368, 341
- Soker, N. & David, L. P. 2003, ApJ, 589, 770
- Soker, N., White, R. E., David, L. P., & McNamara, B. R. 2001, ApJ, 549, 832
- Storchi-Bergmann, T., Nemmen da Silva, R., Eracleous, M., Halpern, J. P., Wilson, A. S., Filippenko, A. V., Ruiz, M. T., Smith, R. C., & Nagar, N. M. 2003, ApJ, 598, 956
- Tamura, T., Kaastra, J. S., Peterson, J. R., Paerels, F. B. S., Mittaz, J. P. D., Trudolyubov, S. P., Stewart, G., Fabian, A. C., Mushotzky, R. F., Lumb, D. H., & Ikebe, Y. 2001, A&A, 365, L87
- Tucker, W. & David, L. P. 1997, ApJ, 484, 602
- Venturi, T., Dallacasa, D., & Stefanachi, F. 2004, A&A, 422, 515

Wise, M. W., McNamara, B. R., & Murray, S. S. 2004, *ApJ*, 601, 184

Young, A. J., Wilson, A. S., & Mundell, C. G. 2002, *ApJ*, 579, 560